

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

### **Long-Term Fine Particulate Matter Exposure and Major Depressive Disorder in a Community-Based Urban Cohort**

Kyoung-Nam Kim, Youn-Hee Lim, Hyun Joo Bae, Myounghee Kim, Kweon Jung, and Yun-Chul Hong

#### **Table of Contents**

**Table S1.** Descriptive characteristics of the study participants included and excluded in the analyses

**Table S2.** Distribution of the study participants (N = 27,270) and the annual PM<sub>2.5</sub> concentration ( $\mu\text{g}/\text{m}^3$ ) in 2007 by districts in Seoul, Republic of Korea

**Table S3.** Hazard ratios of major depressive disorder for an increase of 10  $\mu\text{g}/\text{m}^3$  in the annual PM<sub>2.5</sub> concentration in 2007, stratified by sex and age

**Table S4.** Hazard ratios of major depressive disorder for an increase of 10  $\mu\text{g}/\text{m}^3$  in the annual PM<sub>2.5</sub> concentration in 2007, stratified by household income level

**Table S5.** Hazard ratios of major depressive disorder for an increase of 10  $\mu\text{g}/\text{m}^3$  in the moving average PM<sub>2.5</sub> concentration with a random effect of district

**Table S6.** Hazard ratios of depressive disorder for an increase of 10  $\mu\text{g}/\text{m}^3$  in the PM<sub>2.5</sub> concentration, stratified by different definitions of outcome

**Table S7.** Hazard ratios of major depressive disorder for an increase of 10  $\mu\text{g}/\text{m}^3$  in the PM<sub>2.5</sub> concentration after excluding or including participants who had major depressive disorder until December 31, 2007

**Table S8.** Hazard ratios of major depressive disorder for an increase of 10  $\mu\text{g}/\text{m}^3$  in the PM<sub>2.5</sub> concentration after excluding or including participants who died from intentional self-injury

**Table S9.** Hazard ratios of major depressive disorder for an increase of  $10 \mu\text{g}/\text{m}^3$  in the  $\text{PM}_{2.5}$  concentration in the study population including those without health examination results

**Figure S1.** Flow chart of individuals who met the inclusion/exclusion criteria.

**Figure S2.** The annual average  $\text{PM}_{2.5}$  concentrations between 2007 and 2010 in the 25 districts in Seoul.

**Figure S3.** Penalized regression spline model of the average  $\text{PM}_{2.5}$  concentration in 2007 with a log-transformed hazard ratio. Solid lines, spline curve; shaded area, 95% confidence interval. The model is adjusted for sex, age (5-year group), household income, smoking status, alcohol consumption, regular exercise, the size of the population, proportion of the married person among population aged 15 years or older, economic and social environment satisfaction, the number of clients of the social welfare facilities per capita, and deprivation index in the district in which each participant resided.